

Diane Walder, MD Stacy Chimento, MD Lucy Chen, MD 305.866.2177 (office) 305.866.5302 (fax)

Picoway Laser Pre & Post Procedure Instructions

Precautions to take before your Picoway laser treatment

- No sun exposure, tanning beds and sunless tanning cream for 4 weeks prior to treatment. Sun exposure decreases the effectiveness of the laser treatment and can increase the chance of post treatment complications.
- Avoid any irritants to your skin, such as any products containing Tretinoin, Retinol, Benzoyl Peroxide, Glycolic/Salicylic Acids, Astringents or Vitamin C for one week before your treatment.
- Use broad spectrum UVA/UVB sunscreen with a SPF of 30 or higher. Apply to the treated area every 2 hours when exposed to the sun and it is recommended to make this a part of your skin care routine.
- Remove all makeup, creams or oils prior to treatment.

Instructions following your laser treatment: General (Pigment and Skin Rejuvenation):

- Cleanse the treated area at least daily with water and mild soap, and them pat the area dry.
- Do not rub or scratch the treated area
- If crusting/scabbing occurs, do not shave or pick area. Apply Aquaphor ointment (tattoo) or other moisturizer (face) to the are 2-3 times a day. Keep the area moist, and let the crusting/scabbing resolve on its own.
- Avoid any irritants to your skin, such as any products containing Tretinoin, Retinol, Benzoyl Peroxide, Glycolic/Salicylic Acids, Astringents or Vitamin C for one week after your treatment.
- Discomfort may be relieved by cold gel packs and/or an over the counter pain reliver, such as acetaminophen.
- Avoid contact sports or any other activity that could cause injury of the treated area.
- Avoid swimming, soaking or using hot tubs/whirlpools until the skin heals.
- Contact physician if there is any indication of infection (redness, tenderness or pus)

Tattoo:

- After cleaning and while skin is still moist, apply a thin layer of Aquaphor ointment to the treated tattoo.
- Apply a non-stick pad over the tattoo until it is healed.
- Avoid sun exposure to the treated area. Uses a broad-spectrum UVA/UVB sunscreen with an SPF of 30 or higher. Apply to the treated area every 2 hours when exposed to the sun and it is recommended to make this part of your skin care routine.
- Clean area daily with mild soap and water and pat dry.
- Do not rub or scratch the area.



Diane Walder, MD Stacy Chimento, MD Lucy Chen, MD 305.866.2177 (office) 305.866.5302 (fax)

- Discomfort may be relieved by using cool gel packs or acetaminophen.
- If blistering occurs, keep the area moist by applying Aquaphor 3 times per day or antibiotic ointment per recommendation of the physician. Do not enter swimming pools or hot tubs until treated areas are healed.
- No swimming or using hot tubs for 48 hours post treatment.